SEP 3 0 2002

RECEIVED

OCT 0 5 2002

FANEMARK		TECH CENT	EP21600/290
Form PTO-1449			
ATTY DOCKET NO. 85-02	SERIAL NO. 10/049,137	FILING DATE June 3, 2002	
APPLICANT Ogas and Somerville		GROUP 1653	

U.S. PATENT DOCUMENTS

,	U.S. PATENT DOCUMENTS							
Exmr. Initial		Document Number	Date	Name		Class	Subclass	Filing Date if Appropriate
			FOREIGN	PATENT DOCUM	ENTS			
		Document Number	Date	Country	Class	s	Subclass	Translation Yes/No
	0	THER PRIOR	ART (inclu	ding Author, Title, De	ate, Pe	ertinent	Pages, etc.)
Am		Alonso-Blanco, C. et al., "Development of an AFLP based linkage map of Ler, Col and Cvi Arabidopsis thaliana ecotypes and construction of a Ler/Cvi recombinant inbred line population," (Apr. 1998) The Plant Journal 14(2): 259-271				vi recombinant		
		, I' -	Blázquez, M.A. et al., "LEAFY expression and flower initiation in Arabidosis," (1997) Development 124: 3835-3844				Arabidosis,"	
			Blázquez, M.A. et al., "Gibberellins promote flowering of Arabidopsis by activating the LEAFY promoter," (May 1998) <i>The Plant Cell</i> 10: 791-800					
		1 1	- Bruggemann, E. et al., "Analysis of fast neutron-generated mutants at the Arabidopsis thaliana HY4 locus," (1996) The Plant Journal 10: 755-760					
		1 1	Cowell, I.G. and Austin, C.A., "Self-association of chromo domain peptides," (1997) Biochimica et Biophysica Acta 1337: 198-206					
		chron	Delmas, V. et al., "A mammalian DNA-binding protein that contains a chromodomain and an SNF2/SWI2-like helicase domain," (March 1993) <i>Proc. Natl. Acad. Sci.</i> 90: 2414-2418					
		Eisen distin	Eisen, J.A. et al., "Evolution of the SNF2 family of proteins: subfamilies with distinct sequences and functions," (1995) Nucleic Acis Research 23(14): 2715-2723					
		and re	Finkelstein, R.R. and Zeevaart, J.A.D., "Gibberellin and abscisi acid biosynthesis and response," (1994) <i>Arabidopsis</i> E.M. Meyerowitz and C.R. Somerville, Cold Spring Harbor Laboratory Press: 523-553					
			Goldberg, R.B. et al., "Plant embryogenesis: Zygote to seed," (Oct. 1994) Science 266: 605-614			t. 1994) <i>Science</i>		
M		Grunstein, M., "Histone acetylation in chromatin structure and transcription," (Sept. 1997) Nature 389: 349-352			nscription," (Sept.			

Examiner: Ab & A

Date: 6/20/05

Sheet 2 of 4

Form PTO-1449

RECEIVED

ATTY DOCKET NO. 85-02 SERIAL NO. 10/049,137 FILING DATE June 3, 2002 OCT 0 5 2002

APPLICANT Ogas and Somerville GROUP 1653

TECH CENTER 1600/2900

AM	Hirschhorn, J.N. et al., "Evidence that SNF2/SWI2 and SNF5 activate transcription in yeast by altering chromatin structure," (1992) Genes & Development 6: 2288-2298
	Hooley, R., "Gibberellins: perception, transduction and responses," (1994) Plant Molecular Biology 26: 1529-1555
	Huttly, A.K. and Phillips, A.L., "Gibberellin-regulated plant genes," (1995) Physiologia Plantarum 95: 310-317
	Jin, Y.H. et al., "Isolation and characterization of hrp1 ⁺ , a new member of the SNF2/SWI2 gene family from the fission yeast Schizosaccharomyces pombe," (Feb. 1998) <i>Mol. Gen. Genet.</i> 257: 319-329
	Kehle, J. et al., "dMi-2, a hunchback-interacting protein that functions in polycomb repression," (Dec. 1998) Science 282: 1897-1900
	Khavari, P.A. et al., "BRG1 contains a conserved domain of the SWI2/SNF2 family necessary for normal mitotic growth and transcription," (Nov. 1993) Nature 366: 170-174
	Koonin, E.V. et al., "The chromo superfamily: new members, duplication of the chromo domain and possible role in delivering transcription regulators to chromatin," (1995) Nucleic Acids Research 23: 4229-4233
	Koornneef, M. et al., "A gibberellin insensitive mutant of Arabidopsis thaliana," (1985) Physiol. Plant 65: 33-39
	Koornneef, M. et al., "The isolation of abscisic acid (ABA) deficient mutants by selection of induced revertants in non-germinating gibberellin sensitive lines of Arabidopsis thaliana (L.) Heynh," (1982) Theor. Appl. Genet. 61: 385-393
	Kruger, W. et al., "Amino acid substitutions in the structured domains of histones H3 and H4 partially relieve the requirement of the yeast SWI/SNF complex for transcription," (1995) Genes & Development 9: 2770-2779
	Kwon, H. et al., "Nucleosome disruption and enhancement of activator binding by a human SW1/SNF complex," (Aug. 1994) Nature 370: 477-481
	Logie, C. and Peterson, C.L., "Catalytic activity of the yeast SWI/SNF complex on reconstituted nucleosome arrays," (1997) The EMBO Journal 16: 6772-6782
	Lotan, T. et al., "Arabidopsis LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells," (June 1998) Cell 93: 1195-1205
	McCarty, D.R., "Genetic control and integration of maturation and germination pathways in seed development," (1995) Annu. Rev. Plant Physiol. Plant Mol. Biol. 46: 71-93
	Meinke, D.W., "A homoeotic mutant of Arabidopsis thaliana with leafy cotyledons," (Dec. 1992) Science 258: 1647-1650
	Meinke, D.W. et al., "Leafy cotyledon mutants of arabidopsis," (Aug. 1994) The Plant Cell 6: 1049-1064
Y	Ogas, J. et al., "Cellular differentiation regulated by gibberellin in the Arabidopsis thaliana pickle mutant," (July 1997) Science 277: 91-94
em	Ogas, J. et al., "PICKLE is a CHD3 chromatin-remodeling factor that regulates the transition from embryonic to vegetative development in Arabidopsis," (Nov. 1999) <i>Proc. Natl. Acad. Sci.</i> 96: 13839-13844

Examiner: AGA

Date: 6/20/05

SEP 3 0 2000 SEP 3 1 RADEWARTS

Sheet 3 of 4

Form PTO-1449			RECE	VED
ATTY DOCKET NO. 85-02	SERIAL: NO. 10/049,137	FILING DATE June 3, 2002		
APPLICANT Ogas and Somerville		GROUP 1653	007 0	2002

		ሰባበ
	TECH CENTER 1600/2	300
Am	Parcy, F. et al., "The ABSCISIC ACID-INSENSITIVE3, FUCA3, and LEAFY COTYLEDON1 loci act in concert to control multiple aspects of Arabidopsis seed development," (Aug. 1997) The Plant Cell 9: 1265-1277	
	Peng, J. and Harberd, N.P., "Derivative alleles of the Arabidopsis Gibberellin- Insensitive (gai) mutation confer a wild-type phenotype," (March 1993) The Plant Cell 5: 351-360	
	Prelich, G. and Winston, F., "Mutations that suppress the deletion of an upstream activating sequence in yeast: Involvement of a protein kinase and histone H3 in repressing transcription in vivo," (Nov. 1993) Genetics 135: 665-676	
	Ritchie, S. and Gilroy, S., "Tansley review no. 100 Gibberellins: regulating genes and germination," (Nov. 1998) New Phytol. 140: 363-383	
	Roesler, K.R. et al., "Structure and expression of an Arabidopsis acetyl-coenzyme A carboxylase gene," (1994) Plant Physiol. 105: 611-617	
	Stokes, D.G. et al., "CHD1 is concentrated in interbands and puffed regions of Drosophila polytene chromosomes," (July 1996) Proc. Natl. Acad. Sci. 93: 7137-7142	
	Struhl, K., "Histone acetylation and transcriptional regulatory mechanisms." (Mar. 1998) Genes and Development 12: 599-606	
	Swain, S.M. and Olszewski, N.E., "Genetic analysis of gibberellin signal transduction," (1996) Plant Physiol. 112: 11-17	
	Talon, M., et al., "Accumulation of C ₁₉ -gibberellins in the gibberellin-insenstive dwarf mutant gai of Arabidopsis thaliana (L.) Heynh," (1990) Planta 182: 501-505	
	Tong, J.K. et al., "Chromatin deacetylation by an ATP-dependent nucleosome remodeling complex," (Oct. 1998) Nature 395: 917-921	
	Turner, B.M., "Histone acetylation and control of gene expression," (1991) Journal of Cell Science 99: 13-20	
	Wang, M.L., et al., "Construction of an ≈2 Mb contig in the region around 80 cM of Arabidopsis thaliana chromosome 2," (1997) The Plant Journal 12(3): 711-730	
	West, M.A.L. et al., "Embryogenesis in higher plants: An overview," (Oct. 1993) The Plant Cell 5: 1361-1369	
	West, M.A.L. et al., "LEAFY COTYLEDON1 is an essential regulator of late embryogenesis and cotyledon identity in Arabidopsis," (Dec. 1994) The Plant Cell 6: 1731-1745	
	Wilson, R.N. et al., "Gibberellin is required for flowering in Arabidopsis thaliana under short days," (1992) Plant Physiol. 100: 403-408	
	Wilson, R.N. and Somerville, C.R., "Phenotypic suppression of the gibberelin- insensitive mutant (gai) of Arabidopsis," (1995) Plant Physiol. 108:495-502	
	Woodage, T. et al., "Characterization of the CHD family of proteins," (Oct. 1997) Proc. Natl. Acad. Sci. 94: 11472-11477	
	Zhang, Y. et al., "The dermatomyositis-specific autoantigen Mi2 is a component of a complex containing histone deacetylase and nucleosome remodeling activities," (Oct. 1998) Cell 95: 279-289	
/YM	GenBank Accession Number CAB40760 (Mar. 15, 2000) putative protein [Arabidopsis thaliana]	

Examiner &

Date: 6/20/05

SP 3 0 2000 PE

Sheet 4 of 4
RECEIVED

ATTY DOCKET NO. 85-02 SERIAL NO. 10/049,137 FILING DATE June 3, 2002 OCT 0 5
APPLICANT Ogas and Somerville GROUP 1653

TECH CENTER 1600/2900

SM	GenBank Accession Number AAF13875 (Nov. 30, 1999) chromatin remodeling factor CHD3 [Arabidopsis thaliana]
AM	GenBank Accession Number AL031369 (Aug. 27 1999) Arabidosis thaliana DNA chromosome 2, BAC clone F13D4 (ESSAII project)
Am	GenBank Accession Number AC018722 (Apr. 5, 2000) Arabidopsis thaliana chromosome II section 141 of 255 of the complete sequence. Sequence from clones F13D4, T22F11
EXAMINER	DATE CONSIDERED 6/20/05

*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

12/20/89